

Prepared in collaboration with the National Park Service

Projected Sea-Level Rise and High Tide Flooding at Timucuan Ecological and Historic Preserve, Florida



View from observation tower at Round Marsh in Timucuan Ecological and Historic Preserve (TIMU), Florida, in July 2021. Photograph by Hana Thurman, Cherokee Nation System Solutions, under contract to the U.S. Geological Survey.

Overview

Objective

National parks and preserves in the South Atlantic-Gulf Region (<https://www.nps.gov/aboutus/index.htm>) contain valuable coastal habitats such as tidal wetlands and mangrove forests, as well as irreplaceable historic buildings and archeological sites located in low-lying areas. These natural and cultural resources are vulnerable to accelerated sea-level rise (SLR) and escalating high tide flooding (HTF) events. Through a Natural Resources Preservation Program-funded project during 2021–23, the U.S. Geological Survey (USGS), in collaboration with the National Park Service (NPS), estimated the probability of inundation at Timucuan Ecological and Historic Preserve (TIMU) and several other parks under various SLR scenarios and contemporary HTF thresholds. The maps produced for this effort can be used to assess potential habitat change and explore how infrastructure and cultural resources within the park may be exposed to future flooding-related hazards.

Timucuan Ecological and Historic Preserve

Located in Jacksonville, Florida, along the southern Atlantic coast, TIMU covers 145.35 square kilometers of land (that is, area excluding water and aquatic bed wetlands from the National Oceanic and Atmospheric Administration [NOAA] Coastal Change Analysis Program [NOAA, 2019]) (fig. 1). Most of the park is composed of coastal wetlands, specifically salt marsh (NPS, 2020). Because of SLR, the coastal habitats within TIMU and other national parks and preserves in the region are at the forefront of concern. Specifically, these habitats are threatened unless they have the ability to adapt in place (for example, through vertical adjustment; Morris and others, 2002) or migrate to higher ground (for example, through tidal marsh transgression; Enwright and others, 2016). Similarly, SLR exposes important historic, cultural, and recreational sites to risk of damage or loss.



Base from U.S. Geological Survey digital data
Park boundary from National Park Service (2024)
Shaded bathymetry from U.S. Geological Survey, The National Map
Orthoimagery from Esri and its licensors, copyright 2013, 0.5 meter
Universal Transverse Mercator, zone 17 north
North American Datum of 1983

Figure 1. Location of Timucuan Ecological and Historic Preserve (TIMU) within the State of Florida.

Relative Sea-Level Rise

SLR driven by global climate change is an ongoing and intensifying issue affecting low-lying coastal areas around the world (Sweet and others, 2022). Rates of SLR vary regionally because of factors such as the circulation and density of ocean water, rates of vertical land motion, and the location and volume of ice-mass loss from ice sheets and glaciers (Sweet and others, 2022). These localized rates, referred to as “relative SLR,” are differentiated from the global mean rate.

A recent multiagency report generated five potential global SLR scenarios based on the amount of greenhouse gases emitted into the atmosphere and the rate of ice-mass loss (Sweet and others, 2022). The five scenarios, termed “low,” “intermediate-low,” “intermediate,” “intermediate-high,” and “high,” each predict a different magnitude of future global mean SLR. Along with these global SLR scenarios, we used regional and local projections associated with each scenario to reflect relative SLR rates (fig. 2). Additionally, since there is uncertainty in how various natural processes will affect SLR, rates may vary within a given scenario. Whereas the median SLR projection can be represented by a single value, the “likely” SLR can be represented by a range of values (that is, 17th to 83d percentile) (fig. 2).

High Tide Flooding

HTF occurs when spring tides (that is, maximum tides occurring during a full or new moon when the Earth, Moon,

and Sun are aligned), high winds, or storms cause extreme water levels in coastal areas. For this effort, we used HTF thresholds defined by NOAA, which include thresholds designated “minor,” “moderate,” and “major” that vary on the basis of location (that is, latitude and longitude) (Sweet and others, 2022). Minor HTF events can cause disruptions such as road closures (fig. 3), moderate HTF events can cause greater disruptions and possibly damage, and major HTF events can cause destruction and lead to evacuations (NOAA, 2023).

Today, minor HTF events typically occur only a few days each year along the southern Atlantic Coast, and moderate and major HTF events typically occur less than once per year (Sweet and others, 2022). However, these events are exacerbated by relative SLR, and rapid increases in HTF frequency are expected within the next few decades (Thompson and others, 2021). By 2100, minor HTF is projected to occur more often than every other day under the intermediate-low SLR scenario, meaning that the threshold for a flood today could become the high tide level in the future (Sweet and others, 2018).

The increasing frequency and severity of HTF mean that even minor flooding events are no longer just a nuisance but are becoming a serious issue (Thompson and others, 2021). Cumulatively, the effects of many flooding events can be destructive to infrastructure and assets in low-lying areas (Sweet and others, 2018). Tidal flooding represents a contemporary effect of SLR and may be an indicator of challenges to come (Sweet and others, 2022).




Figure 2. Relative sea-level rise (SLR) scenarios for Timucuan Ecological and Historic Preserve (TIMU), Florida, associated with the five global SLR scenarios from Sweet and others (2022). Sea-level change projected in this figure (modified from Thurman and others [2024]) is relative to the year 1992.



Figure 3. Flooding in September 2022 after Hurricane Ian at Fort Caroline National Memorial in Timucuan Ecological and Historic Preserve (TIMU), Florida. Photograph by Steve Kidd, National Park Service, TIMU.

Key Findings

Relative Sea-Level Rise

Inundation probabilities were mapped for TIMU under the intermediate-low and intermediate-high SLR scenarios for the years 2050 and 2100 by using 1992 as the base year (fig. 4). For these scenarios and timesteps, the median SLR projection and likely range of values (that is, 17th to 83d percentile) are 34 centimeters (cm) for the intermediate-low scenario in 2050 (range: 28–40 cm), 68 cm for the intermediate-low scenario in 2100 (range: 58–80 cm), 44 cm for the intermediate-high scenario in 2050 (range: 34–58 cm), and 159 cm for the intermediate-high scenario in 2100 (range: 121–180 cm) (fig. 2) (Thurman and others, 2024).

Under the intermediate-low scenario in 2050, about 81 percent of the park’s nonwater area was classified as likely to be inundated (that is, probability greater than 0.66); under the same scenario, this percentage increased to about 84 percent by 2100 (fig. 4). For the intermediate-high scenario, about 82 percent of the park’s nonwater area was classified as likely to be inundated by 2050 and about 88 percent by 2100 (Thurman and others, 2024).

High Tide Flooding

Inundation probabilities were mapped for TIMU under NOAA’s minor, moderate, and major HTF thresholds (fig. 5). The minor threshold is 56 cm (plus or minus $[\pm]$ 19 cm, the

root mean square error of linear regression, which is the square root of the average of the squared difference between projected values and actual values) above mean higher high water, and the moderate and major thresholds are 84 cm (± 25 cm) and 123 cm (± 39 cm) above mean higher high water, respectively (Sweet and others, 2018, 2022). Under all three HTF thresholds, most of the park’s nonwater area was classified as likely to be inundated. The likely inundation area increased from about 83 percent under the minor threshold to about 86 percent under the major threshold (Thurman and others, 2024).

Applications of Maps

These maps of projected inundation from SLR and HTF can be used to assess current and future impacts of rising water levels to natural habitats, historic and cultural resources, and infrastructure within the park and can provide a resource for management decisions such as implementation of the resist-accept-direct (RAD) framework (Schuurman and others, 2020). Additionally, this mapping approach can be utilized to quantify and depict future habitat change driven by SLR, such as tidal saline wetland migration.

Methods

Data associated with this effort, along with details on the methods for the approximation of park areas that may be inundated from SLR and HTF, can be found in a USGS data release (Thurman and others, 2024).




Figure 4. Sea-level rise (SLR) inundation probabilities for Timucuan Ecological and Historic Preserve (TIMU), Florida, under the SLR scenarios (Sweet and others, 2022) intermediate-low for the years *A*, 2050 and *B*, 2100 and intermediate-high for the years *C*, 2050 and *D*, 2100 (Thurman and others, 2024). Sea-level change projected in this figure is relative to the year 1992. Probability is shown using three bins developed for the Intergovernmental Panel on Climate Change (2022): unlikely, probability less than or equal to 0.33; likely as not, probability greater than 0.33 and less than or equal to 0.66; and likely, probability greater than 0.66. Existing water areas were excluded from the analyses and area calculations; KM, kilometer.




Figure 5. High tide flooding (HTF) inundation probabilities for Timucuan Ecological and Historic Preserve (TIMU), Florida, under the National Oceanic and Atmospheric Administration (Sweet and others, 2022) *A*, minor, *B*, moderate, and *C*, major HTF thresholds (Thurman and others, 2024). Probability is shown using three bins developed for the Intergovernmental Panel on Climate Change (2022): unlikely, probability less than or equal to 0.33; likely as not, probability greater than 0.33 and less than or equal to 0.66; and likely, probability greater than 0.66. Existing water areas were excluded from the analyses and area calculations; KM, kilometer.

Acknowledgments

This Fact Sheet includes the results of research funded by the USGS and the NPS Natural Resources Preservation Program. We thank the NPS South Atlantic-Gulf Region and the staff of TIMU, especially Steve Kidd, for input on SLR scenario choices and William Vervaeke from the NPS Southeast Coast Inventory and Monitoring Network for assistance with this project.

References Cited

- Enwright, N.M., Griffith, K.T., and Osland, M.J., 2016, Barriers to and opportunities for landward migration of coastal wetlands with sea-level rise: *Frontiers in Ecology and the Environment*, v. 14, no. 6, p. 307–316.
- Intergovernmental Panel on Climate Change, 2022, *Climate change 2022—Mitigation of climate change. Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*: Cambridge, U.K., and New York, Cambridge University Press, 2,029 p., accessed February 2, 2024, at <https://doi.org/10.1017/9781009157926>.
- Morris, J.T., Sundareshwar, P.V., Nietch, C.T., Kjerfve, B., and Cahoon, D.R., 2002, Responses of coastal wetlands to rising sea level: *Ecology*, v. 83, no. 10, p. 2869–2877.
- National Oceanic and Atmospheric Administration [NOAA], 2019, C-CAP derived land cover—BETA [view 6 datasets—2015–2017 C-CAP derived 10 meter land cover—BETA]: National Oceanic and Atmospheric Administration dataset, accessed April 2022 at <https://coast.noaa.gov/digitalcoast/data/ccapderived.html>.
- National Oceanic and Atmospheric Administration [NOAA], 2023, What is high tide flooding?: National Oceanic and Atmospheric Administration web page, accessed February 2024 at <https://oceanservice.noaa.gov/facts/high-tide-flooding.html>.
- National Park Service [NPS], 2020, Timucuan Ecological and Historic Preserve, Florida: National Park Service web page, accessed May 2023 at <https://www.nps.gov/timu/index.htm>.
- National Park Service [NPS], 2024, Administrative boundaries of National Park System units—National Geospatial Data Asset (NGDA) NPS national parks dataset: National Park Service dataset, accessed February 2024 at <https://irma.nps.gov/DataStore/Reference/Profile/2224545?inv=True>.
- Schuurman, G.W., Hawkins-Hoffman, C., Cole, D.N., Lawrence, D.J., Morton, J.M., Magness, D.R., Cravens, A.E., Covington, S., O'Malley, R., and Fisichelli, N.A., 2020, Resist-accept-direct (RAD)—A framework for the 21st-century natural resource manager: Fort Collins, Colorado, National Park Service, Natural Resource Report NPS/NRSS/CCRP/NRR—2020/2213, 20 p., accessed July 14, 2023, at <https://doi.org/10.36967/nrr-2283597>.
- Sweet, W.V., Dusek, G., Obeysekera, J., and Marra, J.J., 2018, Patterns and projections of high tide flooding along the U.S. coastline using a common impact threshold: Silver Spring, Maryland, National Oceanic and Atmospheric Administration, National Ocean Service, NOAA Technical Report NOS CO-OPS 086, 56 p., accessed July 14, 2023, at https://tidesandcurrents.noaa.gov/publications/techrpt86_PaP_of_HTFlooding.pdf.
- Sweet, W.V., Hamlington, B.D., Kopp, R.E., Weaver, C.P., Barnard, P.L., Bekaert, D., Brooks, W., Craghan, M., Dusek, G., Frederikse, T., Garner, G., Genz, A.S., Krasting, J.P., Larour, E., Marcy, D., Marra, J.J., Obeysekera, J., Osler, M., Pendleton, M., Roman, D., Schmied, L., Veatch, W., White, K.D., and Zuzak, C., 2022, Global and regional sea level rise scenarios for the United States—Updated mean projections and extreme water level probabilities along U.S. coastlines: Silver Spring, Maryland, National Oceanic and Atmospheric Administration, National Ocean Service, NOAA Technical Report NOS 01, 111 p., accessed July 14, 2023, at <https://aambpublicoceanservice.blob.core.windows.net/oceanserviceprod/hazards/sealevelrise/noaa-nos-techrpt01-global-regional-SLR-scenarios-US.pdf>.
- Thompson, P.R., Widlansky, M.J., Hamlington, B.D., Merrifield, M.A., Marra, J.J., Mitchum, G.T., and Sweet, W., 2021, Rapid increases and extreme months in projections of United States high-tide flooding: *Nature Climate Change*, v. 11, no. 7, p. 584–590, accessed July 14, 2023, at <https://doi.org/10.1038/s41558-021-01077-8>.
- Thurman, H.R., Enwright, N.M., Osland, M.J., Passeri, D.L., Day, R.H., and Simons, B.M., 2024, Sea-level rise and high tide flooding inundation probability and depth statistics in Nassau and Duval Counties, Florida: U.S. Geological Survey data release, <https://doi.org/10.5066/P9K7S6V5>.
- By Hana R. Thurman,¹ Nicholas M. Enwright,² Michael J. Osland,² Davina L. Passeri,² Richard H. Day,² and Bethanie M. Simons¹
- ¹Cherokee Nation System Solutions, under contract to the U.S. Geological Survey
²U.S. Geological Survey

For more information about this publication, contact

Director, [Wetland and Aquatic Research Center](#)
U.S. Geological Survey
700 Cajundome Blvd.
Lafayette, LA 70506–3152

For additional information, visit
<https://www.usgs.gov/centers/wetland-and-aquatic-research-center-warcc>

Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Although this information product, for the most part, is in the public domain, it also may contain copyrighted materials as noted in the text. Permission to reproduce copyrighted items must be secured from the copyright owner.

Publishing support provided by
Lafayette Publishing Service Center

ISSN 2327-6932 (online)
<https://doi.org/10.3133/fs20243008>